



### AMISTAR PRO

### PRODUCT USE

#### SUMMARY OF INFORMATION

**FORMULATION AND ACTIVE INGREDIENT:** AMISTAR PRO is a suspo-emulsion containing 100 g/litre (9.81% w/w) azoxystrobin and 280 g/litre (27.5% w/w) fenpropimorph.

**MAPP NUMBER:** 10513

**PACK SIZE:** 5 litres.

**TARGETS:**

**Wheat:** *Septoria tritici*, *Septoria nodorum*, Yellow Rust, Brown Rust and Ear Diseases – moderate control. Can reduce the severity of Take-all.

**Barley:** Powdery Mildew, Net Blotch, Brown Rust and *Rhynchosporium*. Can reduce the severity of Take-all.

**Oats:** Crown Rust and Powdery Mildew.

**Rye:** Brown Rust, Powdery Mildew and *Rhynchosporium*. Can reduce the severity of Take-all.

**Triticale:** Brown Rust, Powdery Mildew and *Rhynchosporium* (reduction). Can reduce the severity of Take-all.

**WATER VOLUME AND BCPC SPRAY QUALITY:** Apply in at least 200 litres of water per hectare using a MEDIUM quality spray (BCPC) at a pressure of at least 2 bar. In dense crops increase the water volume to 250–300 litres per hectare.

**PRACTICAL NOTES:**

CROPS	MAXIMUM INDIVIDUAL DOSE (LITRES PRODUCT/HA)	MAXIMUM TOTAL DOSE (LITRES PRODUCT/HA/ CROP)	LATEST TIME OF APPLICATION
Wheat and barley	2	4	Before early milk stage (GS73) or 5 weeks before harvest, whichever is sooner
Oats, rye and triticale	2	4	Up to and including watery ripe stage (GS71) or 5 weeks before harvest, whichever is sooner

To reduce the risk of resistance developing in target diseases the total number of applications of products containing QoI fungicides made to any cereal crop must not exceed two.

**RAINFASTNESS:** 2 hours.

**BUFFER ZONE RESTRICTIONS:** None.

**FOR USE ONLY AS:** AN AGRICULTURAL FUNGICIDE



This product label is compliant with the CPA Voluntary Initiative (VI) guidance. NAN2442/2005