

SEED TREATMENT

COMPATIBILITY

No compatibilities available for this product.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

GENERAL INFORMATION

Results from trials have shown BERET MULTI to be well tolerated over a wide range of sowing dates and conditions. However under adverse environmental or soil conditions, the germination capacity of seed may be reduced and therefore seed rates should be increased in accordance with good agricultural practice.

BERET MULTI is effective against benzimidazole resistant and sensitive forms of *Microdochium nivale*.

DISEASES CONTROLLED**Winter and spring wheat**

Microdochium nivale, *Fusarium culmorum* and *Septoria nodorum* resulting in reductions in Seedling Blight, Stem-Base Browning, Snow Mould and *Septoria* Seedling Blight. Seed-borne and soil-borne Common Bunt (*Tilletia caries*). Loose Smut (*Ustilago tritici*).

Winter and spring barley

Microdochium nivale and *Fusarium culmorum* resulting in reductions in Seedling Blight, Stem-Base Browning and Snow Mould.

Covered Smut (*Ustilago hordei*), Loose Smut (*Ustilago nuda*) and Barley Leaf Stripe (*Pyrenophora graminea*).

Winter and spring oats

Microdochium nivale and *Fusarium culmorum* resulting in reductions in Seedling Blight and Stem-Base Browning. Pyrenophora Leaf Spot (*Pyrenophora avenae*).

CROP SPECIFIC INFORMATION**Crops**

BERET MULTI is recommended as a seed treatment for use on winter and spring crops of wheat, barley and oats.

Timing

Before drilling.

Rates of use

2.0 litres BERET MULTI per tonne of seed.

APPLICATION

BERET MULTI should be applied directly to the seed using conventional seed treatment equipment.

Continuous flow treaters should be calibrated using BERET MULTI before use.

Do not apply BERET MULTI to cracked, split or sprouted seed.

STORAGE AFTER TREATMENT

Ensure seed is of a satisfactory moisture content before storage. Keep in a dry place correctly labelled.

S E E D T R E A T M E N T

Seed bag labels are available from Syngenta Crop Protection UK Ltd.

DRILLING

Seed treated with BERET MULTI may affect the flow rate of seed through drills. It is therefore important to check the calibration of the drill with BERET MULTI treated seed before drilling commences.

SEED BAG LABEL

This seed has been treated with BERET MULTI

Contains fludioxonil and flutriafol for control of seed-borne diseases on winter and spring crops of wheat, barley and oats and also soil-borne bunt.

MAPP 13017

PRECAUTIONS

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling treated seed or contaminated surfaces.

THIS SEED HAS BEEN CHEMICALLY TREATED.

DO NOT HANDLE seed unnecessarily.

TREATED SEED MUST NOT BE USED as food or feed.

DO NOT RE-USE SACKS OR CONTAINERS THAT HAVE CONTAINED TREATED SEED for food or feed.

KEEP TREATED SEED SECURE from people, domestic stock/pets and wildlife at all times during storage and use.

HARMFUL TO GAME AND WILDLIFE. Treated seed should not be left on the soil surface. Bury spillages.

Do not contaminate surface water or ditches with chemical or used container.

WASH HANDS AND EXPOSED SKIN before meals and after work.

DO NOT APPLY TREATED SEED FROM THE AIR.

CAUTION

BERET MULTI may reduce the flow rate of seed in a seed drill. Recalibrate drill with BERET MULTI treated seed before drilling.

PROTECT FROM FROST.

SHAKE WELL BEFORE USE (10 litre container only).

MIX THOROUGHLY BEFORE USE (200 and 1000 litre containers only).

Syngenta Crop Protection UK Limited, CPC4, Capital Park, Fulbourn, Cambridge CB21 5XE
Tel: Cambridge (01223) 883400

S E E D T R E A T M E N T

SAFETY PRECAUTIONS

(a) Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following Personal Protective Equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the product, treated seed or contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH HANDS AND EXPOSED SKIN before meals and after work.

(b) Environmental protection

DO NOT CONTAMINATE SURFACE WATERS OR DITCHES with chemical or used container. TREATED SEED MUST NOT TO BE USED as food or feed.

DO NOT APPLY TREATED SEED FROM THE AIR.

DO NOT RE-USE SACKS OR CONTAINERS THAT HAVE CONTAINED TREATED SEED for food or feed.

LABEL TREATED SEED with the appropriate precautions using printed sacks, labels or bag tags supplied.

(c) Storage and disposal – All refillable containers

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

OPEN THE CONTAINER ONLY AS DIRECTED.

DO NOT RINSE OUT THE CONTAINER.

DO NOT RE-USE CONTAINER for any other purpose.

RETURN EMPTY CONTAINER TO SUPPLIER.

(d) Storage and disposal – All non-refillable containers

EMPTY CONTAINER COMPLETELY and dispose of safely.

KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

BERET MULTI



**DANGEROUS
FOR THE
ENVIRONMENT**

Contains 25 g/l (2.34% w/w) fludioxonil and 25 g/l (2.34% w/w) flutriafol as a flowable concentrate for seed treatment

TOXIC TO AQUATIC ORGANISMS, MAY CAUSE LONG-TERM ADVERSE EFFECTS IN THE AQUATIC ENVIRONMENT.

Keep out of reach of children.

Keep away from food, drink and animal feeding stuffs.

When using do not eat, drink or smoke.

This material and its container must be disposed of in a safe way.

Use appropriate containment to avoid environmental contamination.

To avoid risks to man and the environment, comply with the instructions for use

THE (COSHH) CONTROL OF SUBSTANCES HAZARDOUS TO HEALTH REGULATIONS MAY APPLY TO THE USE OF THIS PRODUCT AT WORK

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL SEED TREATMENT

Crops	Maximum individual dose (litres/tonne)	Maximum number of treatments	Latest time of application
Wheat (seed), Barley (seed) and Oats (seed)	2	One per batch	Before drilling

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

Syngenta Crop Protection UK Limited, CPC4, Capital Park, Fulbourn, Cambridge CB21 5XE
Tel: Cambridge (01223) 883400

In case of toxic or transport emergency ring 01484 538444 any time