

## F U N G I C I D E

**DIRECTIONS FOR USE**

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

**GENERAL INFORMATION**

MENARA is an emulsifiable concentrate containing 250 g/l propiconazole and 160 g/l cyproconazole for use as a foliar fungicide on all commercial varieties of winter wheat and winter and spring barley.

Treatment at early stages of disease development will ensure the best results.

**Caution**

The possible development of resistance to MENARA cannot be predicted. Should it occur MENARA may not give satisfactory control and no responsibility can be accepted for any loss incurred.

MENARA applied to winter wheat in the spring at GS30–33 may, on occasion, cause straw shortening. This effect causes no loss of yield and can be beneficial in a potential lodging situation.

**DISEASES CONTROLLED**

MENARA controls Powdery Mildew, Yellow And Brown Rust, Net Blotch, *Septoria* spp., Ear Diseases and *Rhynchosporium*.

Apply at the first sign of disease.

When applied before or during the period of stem extension in the spring (GS30–32), MENARA will give a useful reduction of Eyespot.

**CROP SPECIFIC INFORMATION**

For use on all varieties of winter wheat and winter and spring barley.

Avoid applying MENARA to crops under stress.

**Timing**

For the control of Powdery Mildew, Yellow And Brown Rust, Net Blotch, *Septoria* spp., Ear Diseases and *Rhynchosporium*, apply at the first sign of disease.

For Eyespot when applied before or during the period of stem extension in the spring (GS30–32), MENARA will give a useful reduction.

**Rates of Use**Winter wheat

Apply MENARA at 0.5 l/ha. A maximum of 1.0 l/ha MENARA may be used per crop.

Barley

Apply MENARA at 0.5 l/ha. A maximum of 0.5 l/ha MENARA may be used per crop.

**MIXING AND SPRAYING**

Apply MENARA in 200–400 litres of water per hectare using a tractor mounted sprayer and a medium spray as defined by BCPC. A spray pressure of 2–3 bars is recommended.

Make sure the sprayer is set to give an even application at the correct volume.

Add half the volume of clean water to the spray tank. Turn on the agitation and add the recommended quantity of MENARA. Continue agitation whilst topping up the spray tank and continue agitation during spraying. Wash out all spray equipment thoroughly with water immediately after use. The spray solution must not be left in the tank and should be used as soon as possible after mixing.

PROTECT FROM FROST

## SAFETY PRECAUTIONS

---

**Precautions marked \* are a legal requirement**

**(a) Operator protection**

\* Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

\* WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

\* WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

\* However engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH CONCENTRATE from skin or eyes immediately.

WASH HANDS AND EXPOSED SKIN before meals and after work.

**(b) Environmental protection**

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

**(c) Storage and disposal**

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

F U N G I C I D E

**MENARA**



**HARMFUL**



**DANGEROUS  
FOR THE  
ENVIRONMENT**

Contains 250 g/l propiconazole and 160 g/l cyproconazole and branched alkyl esters C6-C8 as an emulsifiable concentrate.

Contains propiconazole. May cause an allergic reaction.

**POSSIBLE RISK OF HARM TO THE UNBORN CHILD.  
VERY TOXIC TO AQUATIC ORGANISMS, MAY  
CAUSE LONG-TERM ADVERSE EFFECTS IN THE  
AQUATIC ENVIRONMENT.**

Keep out of reach of children.

When using do not eat, drink or smoke.

Keep away from food, drink and animal feeding stuffs.

Wear suitable protective clothing and gloves.

This material and its container must be disposed of in a safe way.

Use appropriate containment to avoid environmental contamination.

**To avoid risks to man and the environment, comply with the instructions for use.**

THE (COSHH) CONTROL OF SUBSTANCES HAZARDOUS TO HEALTH REGULATIONS MAY APPLY TO THE USE OF THIS PRODUCT AT WORK

**IMPORTANT INFORMATION**

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

For use on winter wheat and winter and spring barley

Crops	Maximum individual dose	Maximum total dose	Latest time of application
Winter wheat	0.5 litre product per hectare	1 litre product/ hectare/crop	Before caryopsis watery ripe stage (GS71)
Winter and spring barley	0.5 litre product per hectare	0.5 litre product/ hectare/crop	Before beginning of anthesis (GS60)

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

Syngenta Crop Protection UK Limited, CPC4, Capital Park, Fulbourn, Cambridge CB21 5XE  
Tel: Cambridge (01223) 883400

**In case of toxic or transport emergency ring 01484 538444 any time.**